




















International Patent Application
No. PCT/EP2004/010582
BioNetWorks GmbH
29607P WO/MDmh







New Claims







1. Use of an 11- β -HSD-type 1 and/or type 2 inhibitor or a pharmaceutically acceptable salt thereof, for the manufacture of a pharmaceutical agent for the prevention and/or treatment of inflammation-induced and/or immune-mediated loss of bone and/or cartilage, wherein said use is for the prevention and/or treatment of osteoporosis, postmenopausal osteoporosis, lytic bone metastases, arthritis, juvenile chronic arthritis and/or adjuvant arthritis, infectious diseases, bone loss by cancer, bone loss by HIV, tooth loss, bone marrow inflammation, synovial inflammation, cartilage and/or bone erosion and/or proteoglycan damage.
2. The use according to claim 1 for the prevention and/or treatment of inflammation-induced and/or immune-mediated loss of bone and/or cartilage in a mammal.
3. The use according to claim 2, wherein the mammal is a human.
4. The use according to claim 1, wherein said use is for the prevention and/or treatment of periodontitis and/or arthritis selected from the group consisting of osteoarthritis and/or rheumatoid arthritis.
5. The use according to any one of claims 1 to 4, wherein the 11- β -HSD-type 1 and/or type 2 inhibitor is 18- β -glycyrrhetic acid.
6. The use according to any one of claims 1 to 4, wherein the 11- β -HSD-type 1 and/or type 2 inhibitor is selected from the group consisting of the following formulas:

Compound Name	Structure
Formula 1	
Formula 2	
Formula 3	
Formula 4	
Formula 5	
Formula 6	
Formula 7	


Formula 8	
Formula 9	
Formula 10	
Formula 11	
Formula 12	
Formula 13	

Formula 14	
Formula 15	
Formula 16	
Formula 17	
Formula 18	
Formula 19	

Formula 20	
Formula 21	
Formula 22	
Formula 23	
Formula 24	
Formula 25	

Formula 26	
Formula 27	
Formula 28	
Formula 29	
Formula 30	
Formula 31	

7. The use according to any one of claims 1-4, wherein the 11- β -HSD-type 1 and/or type 2 inhibitor has the structure of formula I:



formula I

wherein R¹ is

a hydrogen,

a linear or branched C₁-C₁₀ alkyl group,

a linear or branched C₁-C₁₀ alkenyl group,

a linear or branched C₁-C₁₀ alkynyl group,

an ester, amino, halo, hydroxy, carbonyl, carboxy, carboxyphenoxy, C₁-C₄ alkoxy, C₁-C₄ alkoxy carbonyl, C₁-C₄ alkyl amino, di-(C₁-C₄-alkyl)amino, cyano, carboxy amide, carboxy-(C₁-C₄-alkyl)amino, carboxy-di(C₁-C₄-alkyl)sulfo, sulfido (C₁-C₄-alkyl), sulfoxido (C₁-C₄-alkyl), sulfono (C₁-C₄-aminoalkyl) or thio group, a saturated or unsaturated, aromatic or heteroaromatic mono- or polycyclic group,

wherein said cyclic group may be mono- or polysubstituted with an ester, amino, halo, hydroxy, C₁-C₄ alkoxy, carboxy, carbonyl, C₁-C₄ alkoxycarbonyl, carboxyphenoxy, C₁-C₄ alkyl amino, di-(C₁-C₄-alkyl)amino, cyano, carboxy amide, carboxy-(C₁-C₄-alkyl)amino, carboxy-di(C₁-C₄-alkyl)amino, sulfo, sulfido (C₁-C₄-alkyl), sulfoxido (C₁-C₄-alkyl), sulfono (C₁-C₄-alkyl), thio, C₁-C₄ alkyl, C₂-C₄ alkenyl or C₂-C₄ alkynyl group;

R² is

a hydrogen, C₁-C₄ alkyl, carbonyl, ester, amino, halo, carbonyl, hydroxy, carboxy, carboxyphenoxy, C₁-C₄ alkoxy, C₁-C₄ alkoxy carbonyl, C₁-C₄ alkyl amino, di-(C₁-C₄-alkyl)amino, cyano, carboxy amide, carboxy-(C₁-C₄-alkyl)

amino, carboxy-di(C₁-C₄-alkyl), sulfo, sulfido (C₁-C₄-alkyl), sulfoxido (C₁-C₄-alkyl), sulfono (C₁-C₄-alkyl) or thio group;

\mathbb{R}^3 is

a hydrogen,

a linear or branched C₁-C₁₀ alkyl group,

a linear or branched C₁-C₁₀ alkenyl group,


a linear or branched C₁-C₁₀ alkynyl group,

an ester, amino, halo, hydroxy, carbonyl, carboxy, carboxyphenoxy, C₁-C₄ alkoxy, C₁-C₄ alkoxy carbonyl, C₁-C₄ alkyl amino, di-(C₁-C₄-alkyl)amino, cyano, carboxy amide, carboxy-(C₁-C₄-alkyl)amino, carboxy-di(C₁-C₄-alkyl)sulfo, sulfido (C₁-C₄-alkyl), sulfoxido (C₁-C₄-alkyl), sulfono (C₁-C₄-aminoalkyl) or thio group, a saturated or unsaturated, aromatic or heteroaromatic mono- or polycyclic group;


wherein the chemical bond from carbon 13 to 14 is saturated or unsaturated;

or a salt or derivative thereof in the form of an individual enantiomer, diastereomer or a mixture thereof.


8. The use according to claim 1, wherein the 11- β -HSD-type 1 and/or type 2 inhibitor is selected from the group consisting of the following formulas:




Formula 14




Formula 24



Formula 25



9. The use according to any one of claims 1-4, wherein the 11- β -HSD-type 1 and/or type 2 inhibitor has the structure of formula II:



formula II

wherein R¹ is

a hydrogen,

a linear or branched C₁-C₁₀ alkyl group,

a linear or branched C₁-C₁₀ alkenyl group,

a linear or branched C₁-C₁₀ alkynyl group,

an ester, amino, halo, hydroxy, carbonyl, carboxy, carboxyphenoxy, C₁-C₄ alkoxy, C₁-C₄ alkoxy carbonyl, C₁-C₄ alkyl amino, di-(C₁-C₄-alkyl)amino, cyano, carboxy amide, carboxy-(C₁-C₄-alkyl)amino, carboxy-di(C₁-C₄-alkyl)sulfo, sulfido (C₁-C₄-alkyl), sulfoxido (C₁-C₄-alkyl), sulfono (C₁-C₄-aminoalkyl), thio group, a saturated or unsaturated, aromatic or heteroaromatic mono- or polycyclic group,

wherein said cyclic group may be mono- or polysubstituted with an ester, amino, halo, hydroxy, C₁-C₄ alkoxy, carbonyl, carboxy, C₁-C₄ alkoxy carbonyl, carboxyphenoxy, C₁-C₄ alkyl amino, di-(C₁-C₄-alkyl)amino, cyano, carboxy amide, carboxy-(C₁-C₄-alkyl)amino, carboxy-di(C₁-C₄-alkyl)amino, sulfo, sulfido (C₁-C₄-alkyl), sulfoxido (C₁-C₄-alkyl), sulfono (C₁-C₄-alkyl), thio, C₁-C₄ alkyl, C₂-C₄ alkenyl or C₂-C₄ alkynyl group;

R² is a hydrogen or C₁-C₄ alkyl,

R³ and R⁴ are each selected from

a hydrogen

a linear or branched C₁-C₁₀ alkyl group,

a linear or branched C₁-C₁₀ alkenyl group,

a linear or branched C₁-C₁₀ alkynyl group,


an ester, amino, halo, hydroxy, carbonyl, carboxy, carboxyphenoxy, C₁-C₄ alkoxy, C₁-C₄ alkoxy carbonyl, C₁-C₄ alkyl amino, di-(C₁-C₄-alkyl)amino, cyano, carboxy amide, carboxy-(C₁-C₄-alkyl)amino, carboxy-di(C₁-C₄-alkyl)sulfo, sulfido (C₁-C₄-alkyl), sulfoxido (C₁-C₄-alkyl), sulfono (C₁-C₄-aminoalkyl), thio group, a saturated or unsaturated, aromatic or heteroaromatic mono- or polycyclic group;

R⁵ is a hydrogen, C₁-C₄ alky, carbonyl, ester, amino, halo, hydroxy, carboxy, carboxyphenoxy, C₁-C₄ alkoxy, C₁-C₄ alkoxy carbonyl, C₁-C₄ alkyl amino, di-(C₁-C₄-alkyl)amino, cyano, carboxy amide, carboxy-(C₁-C₄-alkyl) amino, carboxy-di(C₁-C₄-alkyl), sulfo, sulfido (C₁-C₄-alkyl), sulfoxido (C₁-


C₄-alkyl), sulfono (C₁-C₄-alkyl) or thio group,

wherein the chemical bond from carbon 8 to 9 is saturated or unsaturated;
wherein the chemical bond from carbon 13 to 14 is saturated or unsaturated;
or a salt or derivative thereof in the form of an individual enantiomer, diastereomer or a mixture thereof.

10. The use according to claim 1, wherein the 11- β -HSD-type 1 and/or type 2 inhibitor is:



11. The use according to claim 6, wherein the 11- β -HSD-type 1 and/or type 2 inhibitor is:



12. The use of any one of claims 1 to 11, wherein the pharmaceutical agent comprises at least one 11- β -HSD-type 1 and/or type 2 inhibitor in combination with at least one active ingredient being effective in the

prevention and/or treatment of inflammation-induced and/or immune-mediated loss of bone and/or cartilage.

13. The use according to any one of claims 1 to 12, wherein the pharmaceutical agent is administered in a dose of 5 to 100 mg/kg body weight per day.
14. The use of any one of claims 1 to 13, wherein the pharmaceutical agent is administered orally, sublingually, intravenously, intramuscularly, intraarticularly, intraarterially, intramedullarily, intrathecally, intraventricularly, intraocularly, intracerebrally, intracranially, respiratorally, intratracheally, nasopharyngeally, transdermally, intradermally, subcutaneously, intraperitoneally, intranasally, enterally, topically, via rectal means, via infusion and/or via implant.
15. The use according to claim 14, wherein the pharmaceutical agent is administered orally.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record.**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.